

Inter-arch Tooth Size Relationships among Different Occlusion Groups of Iraqi Population

Nada M Al-Sayagh
BDS, MSc (Assist Prof)

Dept of Pedod, Orthod, and Prev Dentistry
College of Dentistry, University of Mosul

الخلاصة

الأهداف: كان هدف هذه الدراسة هو تحديد العرض الانسي الوحشي للأسنان الدائمة و نسبة حجم الأسنان بين الفكين و مقارنة هذه المتغيرات بين الجنسين وبين سوء الإطباق المختلفة والإطباق الطبيعي للمراهق العراقي في مدينة الموصل. **المواد والطرق:** اخذ ١٤١ قالب جبسي لطلاب المدارس بعمر ١٣-١٦ سنة من العلاقات الإطباقية المختلفة (الصف الأول ذو الإطباق طبيعي، وسوء الإطباق من الصف الثاني (قسم ١ و ٢) والصف الثالث). قيس عرض الأسنان الانسي الوحشي باستعمال ورنية الأسنان. المتوسط والانحراف المعياري حسباً لجميع المتغيرات واجري التحليل الإحصائي باستخدام اختبار الطالب t، تحليل التباين، اختبار مدى تحليل "دنكن" المتعدد المديات ومعامل ارتباط Pearson. **النتائج:** بالرغم من أن الذكور كان عندهم عرض أكبر لأغلب الأسنان من أولئك في الإناث، لكن البعض من هذه المقاييس كانت غير مختلفة خصوصاً في الصف الأول ذو الإطباق الطبيعي، بينما أهم الاختلافات بين الجنسين وجدت في الصف الثاني قسم ١. اظهر الصف الأول للإطباق الطبيعي ميل نحو الأسنان الصغيرة مقارنة بأصناف سوء الإطباق خصوصاً في مجموعة الذكور بينما اظهر الصف الثالث من سوء الإطباق ميل نحو الأسنان الكبيرة مقارنة بأصناف الإطباق الأخرى خصوصاً في مجموعة الإناث. الصف الأول ذو الإطباق الطبيعي امتلك أعلى نسبة للأسنان الأمامية من ذلك في الصف الثالث من سوء الإطباق و أعلى نسبة للأسنان الكلية من ذلك في مجموعات سوء الإطباق في الإناث، بينما في الذكور كانت نسبة الأسنان الكلية اصغر في الصف الثاني قسم ١ من ذلك في الصف الأول ذو الإطباق الطبيعي. لا يوجد اختلاف بين الجنسين لنسب الأسنان في كل أصناف الإطباق ماعدا الصف الثاني قسم ٢ من سوء الإطباق. **الاستنتاجات:** استنتج بان لكل مجتمع علاقات معينة لحجم الأسنان بين الفكين وهناك معيار للعرض الانسي الوحشي لبعض الأسنان خصوصاً في الصف الثاني قسم ١ و هذه النسب قد تكون إحدى العوامل المهمة المسببة لسوء الإطباق، وهكذا فان هذه الدراسة أثبتت الحقيقة بان تحليل بالتون يجب أن يؤخذ بنظر الاعتبار أثناء التشخيص والعلاج التقوي.

ABSTRACT

Aims: The aim of this study was to determine the mesiodistal tooth width of the permanent dentition, interarch tooth size ratios and to compare these variables between genders and among different malocclusion and normal occlusion groups for Iraqi adolescent in Mosul City. **Materials and Methods:** 141 orthodontic models of school students aged 13 – 16 years of different occlusal relationships (class I normal occlusion, class II (division 1 and 2) and class III malocclusion). Mesiodistal width of teeth were measured by using dental vernier. The mean and standard deviation were calculated. Student's t –test, analysis of variance, Duncan's multiple analysis range test and Pearson's correlation coefficient were used for the statistical analysis. **Results:** Although the males had a larger mesiodistal width of most of the teeth than those in the females, but some of these measurements were not significantly different particularly in class I normal occlusion, while the most significant gender differences were found in class II division 1. Class I normal occlusion showed a tendency toward small teeth than the malocclusion groups particularly in males group, while the class III malocclusion showed a tendency toward larger teeth than the other occlusal categories specially in females group. The class I normal occlusion had a higher anterior tooth ratio than that in class III malocclusion and a higher overall tooth ratio than that in the malocclusion groups in females. While in males the overall tooth ratio was smaller in class II division 1 than that in class I normal occlusion. No gender difference for the tooth ratios in all occlusal categories except in class II division 2 malocclusion. **Conclusions:** It was concluded that interarch tooth size relationships are population specific and there is a gender specific for mesiodistal width of some teeth particularly in class II division 1, and these ratios may be one of the important factors in the cause of malocclusion, thus, this study proved the fact that Bolton's analysis should be taken into consideration during orthodontic diagnosis and therapy.

Key words: Tooth width, interarch ratio, malocclusion, normal occlusion.

Al-Sayagh NM. Inter-arch Tooth Size Relationships among Different Occlusion Groups of Iraqi Population. *Al-Rafidain Dent J.* 2010; 10(1): 89-101.

Received: 14/9/2008

Sent to Referees: 17/9/2008

Accepted for Publication: 3/12/2008

INTRODUCTION

A proper balance should exist between

the mesiodistal tooth size of the maxillary and mandibular arches to ensure proper

occlusion⁽¹⁻³⁾. The size mismatch between the maxillary and mandibular dentition can lead to generalized spacing or crowding or deviation from class I occlusion in the posterior region⁽⁴⁻⁶⁾.

There have been several studies suggesting methods of defining and measuring tooth size discrepancies^(7,8), but the best – known study of tooth size disharmony in relation to treatment of malocclusion was by Bolton⁽⁹⁾ who developed two ratios for estimating tooth size discrepancy by measuring the summed mesiodistal widths of the mandibular to maxillary anterior teeth (anterior ratio) and the total widths of all lower to upper teeth from first to first molar (overall or total arch ratio).

Tooth size variations exist among various ethnic groups⁽¹⁰⁻¹⁷⁾, therefore, different diagnostic standards should be established for each racial group in order to provide an effective diagnostic standard.

A comparative study between Jordanians, Iraqi, Yemenites, and Caucasians reported that Jordanians and Iraqi had larger teeth than the other population⁽¹⁸⁾, the later study, however, didn't discuss the differences in the tooth size between different malocclusions.

Various studies have investigated gender^(11,19-23) and malocclusion⁽²⁴⁻³⁵⁾ differences in the intermaxillary tooth ratios. Arja *et al.*,⁽¹⁹⁾ reported some gender differences, but they couldn't demonstrate any differences between class I and II malocclusion. Another study confirmed the gender difference, but also showed the mesiodistal dimensions of upper teeth to be bigger in class I compared with class II (division 1 and 2) and class III. On the other hand, Akyale *et al.*,⁽³¹⁾ demonstrated no significant difference between the two sexes and among the three malocclusion groups. There are no data available about the inter arch tooth size discrepancies among different malocclusion groups for Iraqi population in Mosul city.

The objectives of this study were: 1) to determine mesiodistal tooth width, anterior and overall Bolton ratios in normal occlusion and different malocclusion for Iraqi sample in Mosul City. 2) to compare the mesiodistal width, anterior and overall Bolton ratios between two genders in normal occlusion and different malocclusions

groups. 3) to compare the mesiodistal width, anterior and overall Bolton ratios among normal occlusion and malocclusions groups. 4) to explore if there is might be a correlation between anterior, overall Bolton ratios and mesiodistal width of permanent dentition.

MATERIALS AND METHODS

The samples for this study consisted of 101 Iraqi students with varying malocclusion and 40 students with class I normal occlusion. All subjects were born and living in Mosul city and were between 13 and 16 years of age. After dental classification, the distribution of the sample were as follows: class I normal occlusion (20 male and 20 female); class II division 1 (20 male and 20 female); class II division 2 (16 male and 15 female) and class III (15 male and 15 female).

The inclusion criteria for the subjects were as follows: All permanent teeth had erupted and were present from right first molar through left first molar. No severe mesiodistal and occlusal tooth abrasion. No residual crown or crown – bridge restoration. No tooth deformity. No record of restoration or stripping of incisors and canine teeth.

In addition, the class I normal occlusion had the following criteria: normal occlusion (Angle class I molar and canine relationship). Harmonious overjet and over bite (2 ± 0.5 mm). No crowding or spacing. No transverse discrepancies.

On the dental cast, each tooth from the maxillary and mandibular right first molar to the left first molar was measured at the largest mesiodistal dimensions to the nearest 0.01 mm, using dental vernier (Müncher model, Dentaurum 042 – 751, Germany) and the same examiner made all measurements.

All statistical analyses were performed using the Stastical Package for Social Sciences (SPSS for windows 98, version 10.0 SPSS Inc., Chicago). The mean and standard deviation for each variable in the different groups of malocclusion and class I normal occlusion were calculated. Comparisons between females and males were made for each variable using Student's *t* – test at $p \leq 0.05$. Analysis of variance was used to determine whether significant dif-

ferences existed among the groups. Duncan's multiple range test were done for test the significance differences at $p \leq 0.05$ among different type of malocclusion and normal occlusion groups. The compared variables were mesiodistal tooth widths, the sum of the six anterior teeth in both arches, the sum of the 12 teeth in both arches, the Bolton's anterior and overall ratios.

Pearson's correlation were done for the Bolton anterior and overall ratios with the other variables in different occlusal categories, for more precision in estimating the degree of significance of "r", the value of probability for "r" in correspondence with the sample size was established

and hence we can say whether "r" is significant at $p \leq 0.05$ level or highly significant at $p \leq 0.01$ level.

RESULTS

Table (1) shows the comparison of the mesiodistal tooth width of the maxillary and mandibular permanent dentition between the males and females group in class I normal occlusion. Although the males had greater mesiodistal tooth than females in most of the teeth but the differences were not statistically significant except the mesiodistal width of the lower first molar which was significantly greater in males than that in females.

Table (1): Comparison of mesiodistal width of upper and lower teeth between males & females groups in Class I normal occlusion group.

Maxillary arch								
Tooth	side	Total		Males (N=20)		Females (N=20)		p -alue
		Mean	SD	Mean	SD	Mean	SD	
central incisor	R	8.32	.439	8.34	.442	8.30	.455	.820
	L	8.34	.419	8.39	.493	8.29	.318	.569
Lateral incisor	R	6.37	.376	6.39	.316	6.36	.454	.850
	L	6.36	.426	6.39	.387	6.33	.487	.754
canine	R	7.46	.467	7.45	.517	7.48	.419	.844
	L	7.46	.466	7.44	.489	7.48	.455	.851
First premolar	R	6.59	.391	6.59	.462	6.60	.298	.966
	L	6.70	.373	6.65	.393	6.77	.353	.443
Second premolar	R	6.32	.464	6.27	.476	6.38	.461	.551
	L	6.32	.468	6.23	.465	6.44	.464	.243
First molar	R	9.90	.448	9.97	.364	9.83	.543	.425
	L	9.87	.445	9.95	.473	9.78	.407	.329
Mandibular arch								
central incisor	R	5.10	.266	5.09	.242	5.12	.304	.777
	L	5.12	.335	5.05	.339	5.21	.323	.240
Lateral incisor	R	5.69	.369	5.64	.302	5.68	.443	.765
	L	5.75	.391	5.68	.345	5.77	.442	.572
canine	R	6.46	.345	6.45	.336	6.39	.355	.648
	L	6.59	.364	6.54	.299	6.59	.448	.723
First premolar	R	6.84	.479	6.82	.492	6.88	.485	.774
	L	6.85	.374	6.89	.409	6.83	.339	.678
Second premolar	R	6.79	.407	6.70	.439	6.90	.348	.192
	L	6.75	.400	6.74	.409	6.78	.393	.783
First molar	R	10.68	.507	10.90	.424	10.09	1.14	.009*
	L	10.72	.579	10.92	.551	10.13	1.29	.045*

* Significant difference at $p \leq 0.05$; **All measurements in millimeter.

Tables (2) and (3) demonstrates the comparison of the mesiodistal tooth width between males and females in class II (division 1 and 2) and class III malocclusion for upper and lower arch respectively. In class II division 1, the males had greater mesiodistal width of most of the teeth than those in females except the upper first premolar, upper left lateral incisor, second premolar, lower left lateral incisor and first

molar.

While in class II division 2, the male had greater mesiodistal width in the upper central incisor and upper right canine, lower first molar, lower right central incisor, lower right canine and lower left lateral incisor. In class III malocclusion, the only significant difference between males and females was found in the upper left canine and premolar.

Table (2): Comparison of mesiodistal width for upper teeth between males & females in different types of malocclusions.

Tooth**	Side	Gender•	Class II div.1			Class II div.2			Class III		
			Mean	SD	P *value	Mean	SD	P *value	Mean	SD	P *value
Central incisor	R	M	8.72	.555	.010*	8.83	.578	.023	9.04	.514	.751
		F	8.18	.499		8.17	.520	*	8.96	.475	
	L	M	8.68	.593	.017*	8.95	.568	.018	9.01	.498	.932
		F	8.17	.479		8.24	.576	*	9.03	.348	
Lateral incisor	R	M	6.95	.560	.032*	6.84	.754	.785	7.16	.461	.302
		F	6.41	.719		6.75	.438		6.85	.671	
	L	M	6.77	.519	.143	6.77	.498	.940	6.88	.561	.708
		F	6.41	.737		6.75	.423		6.76	.751	
Canine	R	M	8.09	.609	.014*	8.09	.512	.027	8.19	.418	.339
		F	7.52	.561		7.53	.454	*	7.92	.628	
	L	M	8.07	.383	.004*	8.06	.517	.079	8.16	.285	.045*
		F	7.50	.580		7.62	.439		7.78	.399	
First pre-molar	R	M	6.94	.408	.103	6.91	.496	.682	6.97	.508	.300
		F	6.67	.420		6.81	.505		7.27	.597	
	L	M	6.93	.262	.181	6.88	.296	.591	6.76	.472	.047*
		F	6.76	.409		6.98	.479		7.38	.646	
Second premolar	R	M	6.76	.309	.054	6.56	.608	.559	6.46	.460	.188
		F	6.47	.456		6.41	.354		6.82	.584	
	L	M	6.59	.437	.930	6.63	.533	.520	6.50	.338	.318
		F	6.57	.519		6.49	.385		6.70	.429	
First molar	R	M	10.74	.598	.047*	10.21	.722	.600	10.29	.616	.697
		F	10.30	.549		10.02	.757		10.43	.827	
	L	M	8.72	.488	.018*	10.12	.642	.469	10.64	.498	.323
		F	8.18	.434		9.89	.641		10.34	.622	

* Significant difference at $p \leq 0.05$; ** All measurements in millimeter. •Number of males in Class II div.1=20, ClassII div 2=16, ClassIII=15; Number of females in Class II div.1=20, Class II div 2=15, ClassIII=15.

Table (3): Comparison of mesiodistal width for lower teeth between males and females in different types of malocclusions.

Tooth**	Side	Gender •	Class II div.1			Class II div.2			Class III		
			Mean	SD	P * value	Mean	SD	P * value	Mean	SD	p * value
Central incisor	R	M	5.48	.291	.007 *	5.56	.466	.036*	5.48	.369	.700
		F	5.09	.419		5.21	.284		5.31	.337	
	L	M	5.43	.426	.046*	5.56	.478	.081	5.54	.241	
		F	5.08	.475		5.24	.313		5.43	.338	
Lateral incisor	R	M	5.95	.363	.024 *	6.05	.651	.256	6.10	.289	.185
		F	5.60	.437		5.80	.329		5.70	.469	
	L	M	5.92	.484	.117	6.17	.527	.012 *	6.20	.295	
		F	5.62	.491		5.71	.478		5.84	.420	
Canine	R	M	6.96	.339	.000 *	7.03	.317	.041*	7.11	.349	.162
		F	6.39	.424		6.51	.423		6.69	.325	
	L	M	7.02	.365	.000 *	7.10	.316	.066	7.29	.442	
		F	6.30	.293		6.55	.345		6.91	.296	
First premolar	R	M	7.01	.438	.019 *	7.13	.569	.310	7.15	.533	.148
		F	6.68	.518		6.58	.225		7.29	.609	
	L	M	7.02	.348	.005*	7.12	.521	.283	7.34	.399	
		F	6.53	.509		6.71	.356		7.30	.739	
Second premolar	R	M	7.10	.318	.007 *	7.09	.516	.091	7.28	.439	.870
		F	6.68	.458		6.58	.528		7.15	.504	
	L	M	7.11	.297	.021*	7.12	.433	.130	7.27	.508	
		F	6.76	.472		6.73	.500		7.07	.520	
First molar	R	M	11.05	.552	.052	10.85	.568	.035*	11.10	.551	.634
		F	10.63	.571		10.14	.522		11.07	.764	
	L	M	11.09	.598	.155	11.16	.585	.012*	11.19	.584	
		F	10.77	.581		10.19	.473		11.11	.764	

*Significant difference at $p \leq 0.05$; **All measurements in millimeter; • Number of males in Class II div.1=20, ClassII div 2=16, ClassIII=15; Number of females in Class II div.1=20, Class II div 2=15, ClassIII=15.

Tables (4) and (5) shows the comparison of mesiodistal width of the maxillary and mandibular teeth among different malocclusions and normal occlusion groups in the males and females groups respectively.

Generally, the class I normal occlusion had a lower mean for the mesiodistal width of most teeth than the malocclusion groups. While the class III malocclusion showed a higher values for most of the variables particularly in the males group.

Table (4) : Comparison of mesiodistal tooth width among different malocclusion and normal occlusion groups in males.

	Tooth	side	Class I (n=20)		Class II div.1(n=20)		Class II div.2(n=16)		Class III (n=15)		p value
			mean	Duncan	mean	Duncan	Mean	Duncan	mean	Duncan	
Maxillary arch	Central incisor	R	8.34	a	8.72	ab	8.74	ab	9.04	b	.020
		L	8.39	a	8.68	ab	8.91	b	8.99	b	.051
	Lateral incisor	R	6.39	a	6.95	bc	6.68	ab	7.26	c	.001
		L	6.39	a	6.77	ab	6.71	ab	7.04	b	.017
	Canine	R	7.45	a	8.09	b	8.18	b	8.20	b	.001
		L	7.44	a	8.07	b	8.14	b	8.24	b	.000
	First pre-molar	R	6.59	a	6.93	ab	6.89	ab	7.13	b	.057
		L	6.65	a	6.93	a	6.86	a	6.92	a	.184
	Second premolar	R	6.27	a	6.76	b	6.57	ab	6.49	ab	.052
		L	6.23	a	6.59	ab	6.67	b	6.52	ab	.084
First molar	R	9.97	a	10.75	b	10.31	ab	10.40	ab	.005	
	L	9.95	a	10.74	b	10.23	a	10.77	b	.000	
Mandibular arch	Central incisor	R	5.09	a	5.48	b	5.56	b	5.48	b	.002
		L	5.05	a	5.43	b	5.56	b	5.54	b	.006
	Lateral incisor	R	5.64	a	5.95	ab	6.05	b	6.10	b	.028
		L	5.68	a	5.92	ab	6.17	b	6.20	b	.014
	Canine	R	6.45	a	6.96	b	7.03	b	7.11	b	.000
		L	6.54	a	7.02	b	7.10	b	7.29	b	.000
	First pre-molar	R	6.82	a	7.01	a	7.13	a	7.15	a	.315
		L	6.89	a	7.02	ab	7.12	ab	7.34	b	.097
	Second premolar	R	6.70	a	7.10	b	7.09	b	7.28	b	.008
		L	6.74	a	7.11	b	7.12	b	7.27	b	.011
First molar	R	10.90	a	11.05	a	10.85	a	11.10	a	.625	
	L	10.92	a	11.09	a	11.16	a	11.19	a	.655	

*Different letters horizontally mean significant difference at $p \leq 0.05$. All measurements in millimeter.

Table (5) : Comparison of mesiodistal tooth width among different malocclusion and normal occlusion groups in females group.

	Tooth	side	Class I (n=20)		Class II div.1(n=20)		Class II div.2(n=15)		Class III (n=15)		P value
			mean	Duncan	mean	Duncan	Mean	Duncan	mean	Duncan	
Maxillary arch	Central incisor	R	8.30	A	8.18	a	8.36	a	8.86	b	.040
		L	8.29	A	8.17	a	8.40	a	8.93	b	.005
	Lateral incisor	R	6.36	A	6.14	a	6.88	a	6.80	a	.167
		L	6.33	A	6.41	a	6.75	a	6.68	a	.384
	Canine	R	7.48	Ab	7.52	ab	7.43	a	7.93	b	.154
		L	7.48	A	7.50	a	7.48	a	7.78	a	.424
	First premolar	R	6.60	A	6.67	a	6.73	a	7.17	b	.028
		L	6.77	A	6.76	a	6.91	ab	7.27	b	.060
	Second premolar	R	6.38	Ab	6.47	ab	6.36	a	6.80	b	.156
		L	6.44	A	6.58	a	6.44	a	6.68	a	.594
First molar	R	9.83	A	10.30	a	9.85	a	10.37	a	.109	
	L	9.77	A	10.30	b	9.70	a	10.27	b	.012	
Mandibular arch	Central incisor	R	5.12	A	5.09	a	5.21	a	5.31	a	.000
		L	5.21	a	5.08	a	5.24	a	5.43	a	.000
	Lateral incisor	R	5.74	A	5.60	a	5.80	a	5.70	a	.001
		L	5.83	A	5.62	a	5.71	a	5.84	a	.012
	Canine	R	6.46	A	6.39	a	6.51	a	6.69	a	.000
		L	6.65	Bc	6.30	a	6.55	ab	6.91	c	.000
	First premolar	R	6.88	Ab	6.68	a	6.58	a	7.29	b	.003
		L	6.81	A	6.53	a	6.71	a	7.30	b	.014
	Second premolar	R	6.90	Ab	6.68	a	6.58	a	7.15	b	.088
		L	6.75	A	6.76	a	6.73	a	7.07	a	.404
First molar	R	10.41	A	10.63	ab	10.14	a	11.07	b	.000	
	L	10.47	Ab	10.77	bc	10.19	a	11.11	c	.000	

*Different letters horizontally mean significant difference at $p \leq 0.05$. All measurements in millimeter.

As shown in Table (6), anterior teeth ratio was not significantly different among different malocclusion and normal occlusion group in males. While in females group, the class I normal occlusion had a higher value than that in class III. The overall ratio was significantly smaller in class II division 1 when compared with class I normal occlusion in males. While in females group, the class I normal occlu-

sion had a higher value than that in malocclusion groups. The males had a higher overall teeth ratio when compared with females in class II division 2 as demonstrated in Table (7), while the other types of malocclusion and normal occlusion showed no significant difference in the anterior and overall teeth ratio between males and females.

Table (6) : Comparison of Bolton ratios among normal occlusion and malocclusion groups in males and females.

Variable	Type of occlusion	Males				Female			
		Mean	SD	P value	Duncan	Mean	SD	P value	Duncan
Sum of upper anterior teeth	CI I	44.39	2.23	.001	A	44.24	2.08	.000	A
	CI II 1	47.27	2.78		B	44.19	3.02		A
	CI II 2	47.35	2.33		B	45.30	1.78		AB
	CI III	48.77	2.46		B	46.98	2.62		B
Sum of lower anterior teeth	CI I	34.45	1.39	.000	A	35.00	1.95	.201	A
	CI II 1	36.76	1.74		B	34.08	1.99		A
	CI II 2	37.47	2.08		B	35.03	1.96		A
	CI III	37.72	1.47		B	35.88	1.73		A
Sum of upper 12 teeth	CI I	90.05	4.23	.000	A	90.03	3.55	.000	A
	CI II 1	95.97	3.95		B	91.28	4.88		A
	CI II 2	94.88	4.47		B	91.29	4.03		A
	CI III	96.99	3.47		B	95.54	4.87		B
Sum of lower 12 teeth	CI I	83.42	3.27	.001	A	83.23	3.09	.050	A
	CI II 1	87.26	2.96		B	82.12	4.34		A
	CI II 2	87.82	4.03		B	81.96	4.47		A
	CI III	89.05	3.74		B	86.87	4.79		B
Bolton anterior ratio	CI I	77.70	2.80	.523	A	79.15	3.01	.096	B
	CI II 1	77.85	2.68		A	77.20	2.71		AB
	CI II 2	79.15	2.35		A	77.32	2.56		AB
	CI III	77.39	1.88		A	76.30	2.24		A
Bolton overall ratio	CI I	92.68	1.72	.061	B	92.47	1.71	.001	B
	CI II 1	90.96	1.65		A	89.99	1.83		A
	CI II 2	92.58	2.02		AB	89.78	1.93		A
	CI III	91.82	1.83		AB	90.85	1.38		A

*Vertically for each variable means with the different letter are significantly different at $p \leq 0.0$; Number of males in Class I=20, Class II div.1=20, ClassII div 2=16, ClassIII=15; Number of females in Class I=20, Class II div.1=20, Class II div 2=15, ClassIII=15.

Table (7) : Comparison of Bolton ratio between males and females in normal occlusion and malocclusion groups.

	Type of occlusion	Males		Females		p value
		Mean	SD	Mean	SD	
Sum of upper anterior teeth	CI I	44.39	2.23	44.24	2.08	.864
	CI II 1	47.27	2.78	44.19	3.02	.008
	CI II 2	47.35	2.33	45.30	1.78	.025
	CI III	48.77	2.46	46.98	2.62	.385
Sum of lower anterior teeth	CI I	34.45	1.39	35.00	1.95	.395
	CI II 1	36.76	1.74	34.08	1.99	.001
	CI II 2	37.47	2.08	35.03	1.96	.020
	CI III	37.72	1.47	35.88	1.73	.200
Sum of upper 12 teeth	CI I	90.05	4.23	90.03	3.55	.993
	CI II 1	95.97	3.95	91.28	4.88	.008
	CI II 2	94.88	4.47	91.29	4.03	.138
	CI III	96.99	3.47	95.54	4.87	.927
Sum of lower 12 teeth	CI I	83.42	3.27	83.23	3.09	.881
	CI II 1	87.26	2.96	82.12	4.34	.001
	CI II 2	87.82	4.03	81.96	4.47	.019
	CI III	89.05	3.74	86.87	4.79	.986
Bolton anterior ratio	CI I	77.70	2.80	79.15	3.01	.208
	CI II 1	77.85	2.68	77.20	2.71	.522
	CI II 2	79.15	2.35	77.32	2.56	.318
	CI III	77.39	1.88	76.30	2.24	.510
Bolton overall ratio	CI I	92.68	1.72	92.47	1.71	.747
	CI II 1	90.96	1.65	89.99	1.83	.142
	CI II 2	92.58	2.02	89.78	1.93	.014
	CI III	91.82	1.83	90.85	1.38	.657

* Significant difference at $p \leq 0.05$; Number of males in Class I=20, Class II div.1=20, ClassII div 2=16, ClassIII=15; Number of females in Class I=20, Class II div.1=20, Class II div 2=15, ClassIII=15.

The correlation of the anterior and overall tooth size ratio with the mesiodistal width of the maxillary and mandibular

teeth in normal occlusion and malocclusion groups were presented in Table (8).

Table (8): Correlation between anterior, overall Bolton ratio and mesiodistal width of teeth in normal occlusion and malocclusion groups.

Tooth	Side	CI I		CI II div.1		CI II div.2		CI III			
		Anterior ratio	Overall ratio	Anterior ratio	Overall ratio	Anterior ratio	Overall ratio	Anterior ratio	Overall ratio		
Maxillary arch	Central incisor	R	-.206	-.268	-.290	-.111	-.008	.249	-.483*	-.057	
		L	-.383*	-.416*	-.290	-.164	.150	.251	-.696**	-.242	
	Lateral incisor	R	-.494**	-.499**	-.484**	-.245	.008	-.002	-.371	.091	
		L	-.444	-.307	-.552**	-.263	.009	.063	-.299	.038	
	Canine	R	-.158	-.270	-.270	-.193	.169	.221	-.366	.193	
		L	-.337	-.273	-.372*	-.235	.185	.341	-.076	.272	
	First premolar	R	-.134	-.227	-.273	-.148	.136	-.090	.342	.261	
		L	-.023	-.260	-.303	-.163	.052	-.308	.197	.139	
	Second premolar	R	-.131	-.474	-.147	-.185	-.062	-.173	-.192	-.277	
		L	-.059	-.400*	-.132	-.163	-.119	-.210	-.078	-.444	
	First molar	R	-.086	-.421*	.161	-.049	.361	.087	.046	-.161	
		L	-.056	-.483*	.262	.138	.225	.069	.145	.105	
	Mandibular arch	central incisor	R	.298	.109	.094	.121	.462	.479*	-.275	.321
			L	.057	-.144	.220	.147	.249	.389	-.091	.156
Lateral incisor		R	.589**	.197	.017	.054	.494*	.367	-.155	.149	
		L	.332	-.103	.110	.024	.495*	.496*	.078	.348	
Canine		R	.102	-.088	-.141	-.016	.511*	.476*	.561*	.497*	
		L	.361	-.008	-.098	.053	.583**	.478*	.393	.504*	
First premolar		R	-.230	-.085	-.141	.139	.358	.126	.220	.318	
		L	-.298	-.139	-.085	.280	.341	.368	.076	.316	
Second premolar		R	-.213	-.132	-.174	.127	.314	.319	.110	.382	
		L	-.224	-.165	-.015	.252	.294	.438	.076	.404	
First molar		R	-.333	.088	-.135	.180	.034	.443	-.324	.115	
		L	-.454*	-.084	-.265	.149	.137	.532*	-.367	.164	
Sum of upper anterior teeth			-.404*	-.407*	-.444*	-.233	.119	.266	-.483*	.059	
Sum of upper 12 teeth			-.277	-.481	-.310	-.200	.154	.095	-.221	.003	
Sum of lower anterior teeth			.371*	-.012	.040	.075	.561**	.535*	.136	.440	
Sum of lower 12 teeth			-.070	-.076	-.078	.166	.435	.532*	.012	.400	
Anterior ratio			1.000	.521**	1.000	.639**	1.000	.676**	1.000	.539*	
Overall ratio			.521**	1.000	.639**	1.000	.676**	1.000	.539*	1.000	

* Significant difference at $p \leq 0.05$; **Significant difference at $p \leq 0.01$.

DISCUSSION

Discrepancies in tooth size should be known at the initial diagnosis and treatment planning stages, if perfect results in orthodontic finishing are to be achieved. The treatment alternative for the tooth size discrepancies include restoration of a relatively small teeth, interproximal stripping of a relatively large teeth, modification of crown angulation or inclination and extraction⁽³⁶⁾.

The low standard deviation of values in the Iraqi sample demonstrates low variability, this might be attributed to the strict selection of harmonious cast and to the high accuracy of the vernier calipers (0.01mm) used in this study, this is in agreement with Nourallah *et al.*⁽¹⁴⁾.

The anterior ratio in many studies^(11,12,25,29,38) is some what higher than Bolton ratio⁽⁹⁾, because of greater morphological variability in upper incisor width than that calculated by Bolton on models in patient with an ideal occlusion, this may also be the case in the present study for the anterior inter –arch ratio in class I normal occlusion in females group. Furthermore, the overall ratio in this study in class I normal occlusion for the males and females groups was higher than Bolton ratio⁽⁹⁾. It is relevant to mention the well – known the effect of premolar extractions on the ideal Bolton ratios and is the consequence of the effect on a ratio of reducing the absolute sums of the tooth widths in the same way that the ratio is different for the total arch because lower second premolars are an average, slightly larger than upper premolar⁽¹¹⁾.

The mean values of anterior and overall ratios were not statistically significant between the two genders in class I normal occlusion, class II division 1 and class III malocclusion groups, this confirm the findings of other studies^(14,20,23).

Xia and Wu⁽³⁹⁾ found no significant difference for tooth size ratios between the normal occlusion and malocclusion groups, this confirm the findings of the present study for the anterior ratio in males and also confirm the findings in females except the class III that showed smaller anterior ratio than that in class I normal occlusion, this is disagreement with Nie and Lin⁽²⁵⁾ who demonstrated that a

significant difference was found for intermaxillary tooth size ratios among different malocclusion groups with the ratios showing that class III > class I > class II. In this study, the anterior and overall tooth ratios among the different malocclusions showed no significant differences. This findings was in agreement with other studies^(11,24,30,31,40). Alkofide and Hashim⁽²⁷⁾ also reported no difference in the incidence of tooth size discrepancies among the different malocclusion groups except for the anterior ratio in class III malocclusion.

Although mesiodistal crown width of the most teeth in males group were larger than females. This agree with the previous studies^(14,20,23,25), but some of these measurements were not significantly different particularly in class I normal occlusion.

The most significant gender differences for the mesiodistal teeth width were found in class II division 1. The exact reason for this difference is not well understood. This could be due to sex –linked inheritance and sex –hormonal influence.

Generally, the class I normal occlusion showed a lower mean values for the mesiodistal width of all the teeth than the malocclusion groups, although the difference was not significant in some of measurements in males and females groups. On the other hand, the class III malocclusion showed a higher value for most of the mesiodistal width than other types of malocclusion and normal occlusion group, this is in agreement with Lavelle⁽³⁾ and Xia and Wu⁽³⁹⁾.

The correlational results revealed that the anterior ratio had significant correlation with the sum of the mesiodistal width of the upper anterior teeth in class I normal occlusion, class II division 1 and class III, while in class II division 2, the anterior ratio correlated with the sum of the mesiodistal crown width of the lower anterior teeth, this indicates that the anterior teeth ratio associated with the variation of upper anterior teeth rather than lower anterior teeth in all types of occlusion categories except class II division 2, this support the findings of other studies^(25,37,38) who found that greater morphologic variability in upper incisor width are believed to affect the anterior ratio.

CONCLUSIONS

These findings indicated that population specific standards for interarch tooth size relationships are necessary for clinical assessment. The males showed a tendency of having a significantly larger teeth than females particularly in class II division 1 malocclusion group. Class III malocclusion showed a tendency toward a significantly larger teeth than the other occlusion categories especially in females group. In contrast, class I normal occlusion showed a tendency toward a significantly smaller teeth than the malocclusion groups particularly in males group. The overall tooth ratio in class I normal occlusion was significantly higher than class II division 1 in males and higher than the malocclusion groups in females. In addition, the anterior tooth ratio was significantly higher in class I normal occlusion than in class III malocclusion in females group. No gender difference for the anterior and overall tooth ratio in all occlusal categories except in class II division 2. The anterior tooth ratio associated with the variation of upper anterior teeth.

REFERENCES

1. Thilander B, Rönning O. Introduction to orthodontic. 6th ed. 1998; P: 57.
2. Shellhart WC, Lang DW, Kluemper GT, Hick EP, Kaplan AL. Realability of the Bolton tooth size analysis when applied to crowded dentitions. *Angle Orthod.* 1995; 65: 327–334.
3. Lavelle CLB. Maxillary and mandibular tooth size in different racial groups and in different occlusive categories. *Am J Orthod.* 1972; 61: 29–37.
4. Laino A, Quaremba G, Paduano S, Stanzione S. Prevalence of tooth – size discrepancy among different malocclusion groups. *Prog Orthod* 2002; 4: 37–44.
5. Janosevic M, Filipovic G, Stankovic S, Janjic CT. Influence of the size of incisors on the occurrence of crowding. *Medicine & Biology.* 2006; 13(1): 36–43.
6. Redahan S, Lagerström L. Orthodontic treatment outcome: the relationship between anterior dental relations and anterior inter–arch size discrepancy. *J Orthod.* 2003; 30: 237–244.
7. Kestling HD. The philosophy of the tooth positioning appliance. *Am J Orthod.* 1945; 31: 297–340.
8. Neff CW. Size relationship between the maxillary and mandibular anterior segments of the dental arch. *Angle Orthod.* 1957; 27: 138– 147.
9. Bolton A. Disharmony in tooth size and its relation to the analysis and treatment of malocclusion. *Angle Orthod.* 1958; 28: 113–130.
10. Merz ML, Isaacson RJ, Germane N, Rubenstein LK. Tooth diameters and arch perimeters in a black and a white population. *Am J Orthod. Dentofacial Orthop.* 1991; 100: 53–58.
11. Smith SS, Buschang PH, Watanabe E. Interarch tooth size relationships of 3 populations: "does Bolton's analysis apply?". *Am J Orthod Dentofacial Orthop.* 2000; 117: 169–174.
12. Santoro M, Ayoub ME, Pardi VA, Cangialosi TJ. Mesiodistal crown dimensions and tooth size discrepancy of permanent dentition of Dominican Americans. *Angle Orthod.* 2000; 70: 303–307.
13. Bernabe E, Major PW, Flores-Mir C. Tooth –size ratio discrepancies in a sample of Peruvian adolescents. *Am J Orthod Dentofacial Orthop.* 2004; 125: 3361–3365.
14. Nourallah AW, Splieth CH, Schwahn C, Khurdaji M. Standardizing interarch tooth–size harmony in a Syrian population. *Angle Orthod.* 2005; 75(6): 996–999.
15. Al-Tamimi T, Hashim HA. Bolton tooth–size ratio revisited. *World J Orthod.* 2005; 6(3): 289–295.
16. Pareded V, Gandia JL, Cibrian R. Do Bolton's ratios apply to a Spanish population? *Am J Orthod Dentofacial Orthop.* 2006; 129(3): 428–430.
17. Uysal T, Sari Z. Intermaxillary tooth size discrepancy and mesiodistal crown dimensions for a Turkish population. *Am J Orthod Dentofacial Orthop.* 2005; 128(2): 226–230.
18. Hattab FN, Al-khateeb S, Sultan I. Mesiodistal crown diameters of permanent teeth in Jordanians. *Arch Oral Biol.* 1996; 41: 641– 645.
19. Arya BS, Savara BS, Thomas D, Clarkson Q. Relation of sex and occlusion to mesiodistal tooth size. *Am J Orthod.* 1974; 66: 479–486.
20. Lew KK, Keng SB. Anterior crown

- dimensions and relationship in an ethnic Chinese population with normal occlusion. *Aus Orthod J.* 1991; 12(20): 105–109.
21. Adeyemita Ta, Isiekwe Mc. Comparing permanent tooth size (mesio –distal) of males and females in a Nigerian population. *West Afr J Med.* 2003; 27(3): 219–221.
 22. Schwartz GT, Dean MC. Sexual dimorphism in modern human permanent teeth. *Am J Phy Anthropol.* 2005; 128(2): 312–317.
 23. Singh SP, Goyal A. Mesiodistal crown dimensions of permanent dentition in North Indian children. *J Indian Soc Pedod Prev Dent.* 2006; December, ISSN 0970–4388.
 24. Crosby DR, Alexander CG. The occurrence of tooth size discrepancies among different malocclusion groups. *Am J Orthod Dentofacial Orthop.* 1989; 95: 457–461.
 25. Nie Q, Lin J. Comparison of intermaxillary tooth discrepancies among different malocclusion groups. *Am J Orthod Dentofacial Orthop.* 1999; 116(5): 539–544.
 26. Ta TA, Ling JY, Hagg U. Tooth –size discrepancies among different occlusion groups of Southern Chinese children. *Am. J. Orthod. Dentofacial Orthop.* 2001; 120: 556–558.
 27. Alkofide E, Hashim H. Intermaxillary tooth size discrepancies among different malocclusion classes: a comparative study. *J Clin Pediatr Dent.* 2002; 26: 4383–4387.
 28. Araujo E, Souki M. Bolton anterior tooth size discrepancies among different malocclusion groups. *Angle Orthod.* 2003; 73: 3307–3313.
 29. Kim HS, Shim HY, Nahm DS. A study on the anterior tooth size discrepancies among orthodontic patients with varying malocclusion. *Korean J Orthod.* 2005; 35(6): 420–432.
 30. Al-khateeb SN, Abu Alhaija ES. Tooth size discrepancies and arch parameters among different malocclusion in a Jordanian sample. *Angle Orthod.* 2006; 76(3): 459–465.
 31. Akyalcin S, Dogan S, Dincer B, Eridinc AM, Oncag G. Bolton tooth size discrepancies in skeletal class I individuals presenting with different dental angle classifications. *Angle Orthod.* 2006; 76(4): 637 –643.
 32. Basaran G, Selek M, Hamamc O, Akku Z. Intermaxillary Bolton tooth size discrepancies among different malocclusion groups. *Angle Orthod.* 2005; 76(1): 26–30.
 33. Fattahi HR, Pakshir HR, Hedayati Z. Comparison of tooth size discrepancies among different malocclusion groups. *Eur J Orthod.* 2006; 28(5): 491–495.
 34. Qiong N, Jiuxiang L. Comparison of intermaxillary tooth size discrepancies among different malocclusion groups. *Am J Orthod Dentofacial Orthop.* 1999; 116: 539–544.
 35. Hashim HA, Al-Ghamdi SAF. Tooth width and arch dimensions in normal and malocclusion samples. An odontometric study. *J Contemp Dent Pract.* 2005; 6(2): 036–051.
 36. Fields HW. Orthodontic restorative treatment for relative mandibular anterior excess tooth size problems. *Am J Orthod Dentofacial Orthop.* 1981; 79: 176–183.
 37. Freeman JE, Maskeroni AJ, Lorton L. Frequency of Bolton tooth size discrepancies among orthodontic patients. *Am J Orthod Dentofacial Orthop.* 1996; 110: 24–27.
 38. Heusdens M, Dermaut L, Verbeeck R. The effect of tooth size discrepancy on occlusion: an experimental study. *Am J Orthod Dentofacial Orthop.* 2000; 117(2): 184–191.
 39. Xia Z, Wu XY. The application of dentocclusal measurement in malocclusion. *Stomatology.* 1983; 3: 126–127.
 40. Basaran G, Selek M, Hamaci O, Akku Z. Intermaxillary Bolton tooth size discrepancies among different malocclusion groups. *Angle Orthod.* 2006; 76: 26–30.